

Effects of mineral ions on yeast performance under very high gravity beer fermentation

By

Henry Okwudili, Udeh

Student No: 11626674

MSc Research Project (FST 6099)



University of Venda

A research project submitted in fulfilment of the requirement for the degree of MSc in Food Science and Technology (Course work and mini-dissertation)

Supervisor: Mr. T.E. Kgatla

Co-Supervisor: Professor A.I.O. Jideani

Department of Food Science and Technology, School of Agriculture, University of Venda,
Private Bag X5050, Thohoyandou 0950.

UNIVEN LIBRARY

Library Item : 20141497



March, 2014

ABSTRACT

The effect of three divalent cationic concentrations of Zn^{2+} , Mg^{2+} , and Ba^{2+} on yeast (*Saccharomyces pastorianus*) fermentative performance was investigated at independent and four variable combinations in 24 experimental runs in accordance with the experimental design. Very high gravity wort of 21°P was prepared from barley malt, hops and water, to which the metal and their combinations were supplemented and subsequently pitched using lager brewing strain *S. pastorianus* and allowed for 96 h fermentation. After 96 h of fermentation, highest wort fermentability of %F = 30.16 and %F = 29.53 which were not significantly different ($p \geq 0.05$) were obtained for wort medium supplemented with 12, 900, and 4 ppm (Zn^{2+} , Mg^{2+} , and Ba^{2+}) and 900 and 4 ppm (Mg^{2+} and Ba^{2+}) respectively. Maximum ethanol yield of 8.4347% (v/v) was obtained in wort medium supplemented with the metal combination ratio of 12:900:4 (Zn^{2+} , Mg^{2+} , and Ba^{2+}). For the individual metal concentration levels tested, high wort fermentability of %F = 27.94 and %F = 26.03 which were not significantly different ($p \geq 0.05$) was obtained for wort media supplemented with 700 ppm Mg^{2+} and 2 ppm Ba^{2+} . High ethanol yield of 7.8844% (v/v) and 7.6245% (v/v) was recorded for Mg^{2+} and Ba^{2+} at concentration of 700 ppm and 2 ppm, respectively. Wort pH of 4.75 was found optimal for the yeast activity. Concentrations of 10 ppm Zn^{2+} and 4 ppm Ba^{2+} , were found to reduce yeast performance. Thus, the study established relationships between the metal ions on yeast fermentative performance and the impact of Ba^{2+} on yeast fermentation process – with concentrations of 1 and 2 ppm Ba^{2+} found significant ($p \leq 0.05$).

Keywords: Very high gravity fermentation, brewer's yeast, yeast stress, yeast stress tolerance, yeast performance, mineral ions, fermentability, ethanol yield.