

**THE PREVALENCE OF *Vibrio cholerae* AND OTHER *Vibrio*
spp. IN SURFACE WATER OF RURAL COMMUNITIES IN
THE LIMPOPO PROVINCE OF SOUTH AFRICA**

by

MASINDI WONTONDA

A THESIS SUBMITTED IN FULFILLMENT OF THE REQUIREMENTS OF THE
MASTERS DEGREE IN MICROBIOLOGY (MSc MBY)

to the

DEPARTMENT OF MICROBIOLOGY
SCHOOL OF MATHEMATICAL AND NATURAL SCIENCES
UNIVERSITY OF VENDA
PRIVATE BAG X5050
THOHOYANDOU
0950

Supervisor: Prof N Potgieter

Co-supervisor: Dr AN Traoré

UNIVEN LIBRARY

Library Item : 20172654



April 2016

ABSTRACT

Introduction: *Vibrio cholerae* is a waterborne pathogen that has caused several outbreaks in developing parts of the world due to a lack of access to clean water.

Objectives: The aim of the study was to investigate the prevalence of *Vibrio cholerae* and other *Vibrio* species in surface waters used by rural communities in the Limpopo province of South Africa.

Materials and method: A total of 120 water samples were collected over a period of three months from 13 rivers in the rural communities of the Vhembe district. The study area encompassed 3 of the 4 Vhembe municipalities namely Thulamela, Mutale, Makhado municipalities. The water samples were collected from three points on each river (upstream, middle stream and downstream). Physico-chemical parameters (pH, Temperature, Total dissolved solids (TDS), electric conductivity (EC)) were measured for each sample. Isolation of *Vibrio* spp. was done using membrane filtration technique, followed by pre-enrichment of the bacteria in Alkaline Peptone Water after which the bacteria was cultured onto Thiosulfate Citrate Bile Salts (TCBS) selective agar. Presumptive *Vibrio* isolates were identified using a series of biochemical identification techniques which included: Gram staining, Oxidase test and API 20 E Identification strips. DNA was extracted from all samples which were identified as *Vibrio* spp by the API 20 E strips. Two optimized multiplex conventional PCR assays were used to identify the pathogenic strain of the presumptive *Vibrio* isolates. The first PCR assay targeted 16S rRNA (*Vibrio* species), *ctxA* (cholera toxin) and *sodB* (Multiplex 1) genes and the second PCR assay targeted 16S rRNA *V. cholerae* O1 *rfb* and *V.p flae* (Multiplex 2) genes.

Results and discussion: *Vibrio cholerae* non O1 was the only prevalent strain identified over the study period. Most physico-chemical parameters of the rivers (pH, Temperature, TDS and electric conductivity) were within the (DWA 1998), (WRC 1998) and (WHO 2011) standard guideline limits.